



Basingstoke,
Southampton
& Winchester
District
Prescribing
Committee

Shared Care Guideline for(GP Summary)

It is essential that a transfer of care only takes place with agreement of the GP and when sufficient information has been received. If the GP does not agree to share care they will inform the Consultant responsible for the patient's care.

Specialist Contact Details	Patient ID Label
Name: _____	Surname: _____
Location: _____	Forename: _____
Date: _____	NHS Number: _____
Tel: _____	Date of Birth: _____

Indications

Melatonin is only indicated where non-pharmacological strategies have been tried, but sleep latency and maintenance remains a significant problem. Melatonin may be considered for:

- children and adolescents with chronic fatigue syndrome (CFS) or myalgic encephalomyelitis (ME) who have sleep difficulties in line with NICE CG53 (www.nice.org.uk/guidance/cg53).
- use in children ≥1 year of age with neurodevelopment disability, autism, visual impairment or neuropsychiatric disorders and chronic sleep disturbance, including chronic fatigue syndrome, where both:
 - Symptoms of sleep disturbance have been present for at least six months, or sleep disturbance is so severe that it is causing significant family disturbance.
 - The sleep problem has been thoroughly evaluated, and a disorder of sleep that is likely to respond to melatonin has been diagnosed (e.g. sleep onset insomnia/delayed sleep phase syndrome, likely endogenous melatonin deficiency or phase reversal). Note: importantly disorders of sleep not responsive to melatonin should have been excluded (e.g. limb movement disorder in ADHD and sleep disordered breathing)

Product choice, dose & response

Melatonin is classified as a prescription only medicine in the UK. A number of licensed products are available, although the majority are not licensed for use in children <18 years or for the indications detailed above. For details refer to the BNF or product SmPCs.

The choice of product should be in accordance with General Medical Council guidance on prescribing unlicensed medicines (in [Good Practice in prescribing and managing medicines and devices](#) 2013) and the MHRA guidance on the hierarchy for the use of unlicensed medicines (Appendix 2 [The supply of unlicensed medicinal products](#)).

A licensed product for licensed indication should always be considered first in all cases. But if there is no such product available, or if another product or formulation better suits the patient's needs, the next choice should be a licensed product used "off-label", and finally an unlicensed product (e.g. a UK manufactured "special", a nutritional supplement, or an imported product licensed abroad).

Risks versus benefits should be considered before switching patients already established on an unlicensed or off label product, and it may be more appropriate for them to continue taking that product if it has proved to be effective and well tolerated and they are comfortable with the formulation.

Where there are more than one licensed products for the same cohort of patients, the most cost-effective product should be considered. This applies particularly for use of Slenyto® outside of its licensed indications, i.e. off label (see table below). In this scenario prescribers should consider a more cost effective product for off label use, e.g. Circadin or Melatonin PharmaNord tablets.

The table below provides a summary of melatonin products supported for prescribing locally. A pathway is provided as an Appendix to these guidelines aid formulation choice:

Product	UK product licence	Approved indications	Cost per 30 days ³
Melatonin 1mg, 5mg prolonged-release tablets (Slenyto®)	Yes	Treatment of insomnia in children and adolescents aged 2 to 18 years with autism spectrum disorder (ASD) and/or Smiths-	2mg daily: £41.20 5mg daily: £103.00 10mg daily: £206.00

Prepared by: Kazeem Olalekan, Specialist Pharmacist, Child Health, University Hospital Southampton NHS Foundation Trust and Samantha Owen, Principal Pharmacist, Critical Evaluation and Formulary, University Hospital Southampton NHS Foundation Trust (based on previous guidelines produced by Edward Horne, Pharmacist, Intermediate Care & Community Nursing, Solent NHS Trust, June 2016).
In consultation with: Dr Cathy Hill[and others]
Date prepared: June 2020
Review date: June 2022

		Magenis syndrome where sleep hygiene measures have been insufficient. Data available for up to 2 years' treatment.	
Melatonin 2mg prolonged-release tablets (Circadin®)	Yes	As monotherapy for the short-term (up to 13 weeks) treatment of primary insomnia in adult patients aged ≥ 55 years.	2mg daily: £15.39 6mg daily: £46.17 10mg daily: £76.95
Melatonin PharmaNord® 3mg film-coated tablets ¹ (Prescribe by brand name)	Yes	For short-term treatment (max 5 days) of jet lag in adults (≥18 years) - may be taken for a maximum of 16 treatment periods per year.	3mg daily: £15.95 6mg daily: £31.90 9mg daily: £47.85
Melatonin 1mg in 1ml oral solution (Colonis Pharma Ltd) ²	Yes	For short-term treatment (max 5 days) of jet lag in adults (≥18 years) - may be taken for a maximum of 16 treatment periods per year.	2mg daily: £52.00 5mg daily: £130.00 10mg daily: £260.00

¹ Melatonin PharmaNord 3mg film-coated tablets must be prescribed by brand name. Generic prescriptions for melatonin 3mg tablets, or prescriptions for the Colonis Pharma brand of melatonin 3mg film-coated tablets will incur higher costs.

² Oral solution is less preferred to tablets due to higher cost and content of excipients potentially harmful in children (e.g. propylene glycol, sorbitol).

Other strengths and formulations of liquid melatonin are also available as unlicensed specials (refer to the Drug Tariff Part VIII B). These are not routinely recommended, but may be prescribed in exceptional circumstances when other formulations are not tolerated or suitable. Prices vary but may be lower cost than the licensed oral solution.

³ Prices correct at time of writing but may be subject to change.

Dose

An initial dose of 2 to 3mg once daily is recommended. This may be increased by 2 to 3mg every 7 to 14 days depending on response. Usual maximum dose is 10mg once daily but additional benefits from doses above 6 mg are uncertain.

Administration

- Melatonin should be taken once a day, between 30 to 60 minutes before bedtime.
- Prolonged-release tablets should be taken with or after food, i.e. on a full stomach.
- Immediate release tablets and liquids are best taken on an empty stomach (≥2 hours before or after eating).

Note: split or crushed prolonged-release tablets should be treated as immediate release tablets.

Swallowing difficulties/ enteral feeding tubes

- Leaflets and videos for parents and carers on helping children to take tablets are available on the Medicines for Children website www.medicinesforchildren.org.uk.
- Slenyto® tablets are formulated as a mini tablet, specifically designed as smaller diameter (3mm) to facilitate swallowing in the ASD paediatric population. The SmPC also advises they can be mixed (whole) with food such as yoghurt, orange juice or ice cream and taken immediately.
- The practice of splitting (e.g. halving or quartering) or crushing melatonin tablets (immediate- or prolonged-release) is off label since the manufacturers recommend swallowing the tablets whole. However this practice is well established, particularly with Circadin, and is not associated with safety concerns. The prescription should state if the medication is to be crushed prior to administration. The tablets should be crushed to a fine powder and mixed with water or a small amount of soft food such as jam or yoghurt, or added to 15-30ml of water for administration via enteral feeding tubes (flush with 30ml water before and after giving). Splitting or crushing will result in loss of prolonged-release properties resulting in a faster onset and shorter duration of effect.
- A liquid formulation may also be considered if tablets are not tolerated or unsuitable, but this is generally less preferred due to higher cost, and safety concerns with excipients. Caution is required to ensure the excipient content (e.g. propylene glycol, sorbitol) does not exceed safety limits in children (further information available in PrescQIPP bulletin [245](#)). Unlicensed liquid specials may offer an excipient-free option and may be lower cost than the licensed oral solution (1mg/ml).

Commented [SO1]: Pragmatic – most melatonin products seem to advise on this, including Slenyto and this is what it says in the Medicines for children leaflet. The only one that differs is Circadin but it gets too complicated if you start using different instructions for different brands!

	<p>Response</p> <ul style="list-style-type: none"> Children should start to feel sleepy about 30 minutes after taking a dose of melatonin, possibly slightly longer with a prolonged-release tablet. When swallowed whole, prolonged-release tablets release melatonin over at least 8 hours. Immediate-release tablets or liquids and crushed prolonged-release tablets will not last as long, possibly around 2 to 3 hours. Specialists will review patient's initial response after 4 weeks and continue prescribing until an effective dose is established. A drug holiday should be introduced at least annually to assess the continued need for treatment. Specialists should provide advice to patients, carers and GPs on how to instigate this. It could take place a month before the annual review, with the patient and/or the carer keeping a sleep diary. The outcome of any drug holiday must be recorded in the patient's notes.
Specialist's Responsibilities	<p>Treatment with melatonin should be initiated by an experienced specialist in sleep medicine, CAMHS practitioner, paediatrician or learning disabilities specialist.</p> <ul style="list-style-type: none"> Establish a diagnosis and assess suitability of the patient for treatment, ensuring compatibility of melatonin with concomitant medication. Initiate treatment after discussion of the options available with the patient and carer(s). Discuss benefits and side effects of treatment with the patient/carer including that the medicine or administration method is unlicensed/off label (when applicable) and the lack of data pertaining to potential long-term side-effects of treatment and theoretical risks of delayed sexual maturation. Advise of the need for shared care (once dose stabilised), and obtain appropriate consent to treatment. Write to the GP when the melatonin is initiated, to ask the GP whether he/she is willing to participate in the ongoing prescribing and general care as outlined in this shared care agreement (published online at https://www.westhampshireccg.nhs.uk/shared-care-guidelines). Prescribe the initial supply (for at least 4 weeks treatment) and continue prescribing until the patient is stable on a particular dose and formulation. Once stable, provide the GP with: details of diagnostic information; dose and preparation of melatonin to prescribe (including clinical justification for the choice of preparation), and duration of treatment before specialist review. Offer the patient outpatient appointments at least annually and regular appointments with the community teams or paediatric support team. At these appointments the efficacy and safety of melatonin should be reassessed, and treatment discontinued if appropriate. A drug holiday should be carried out at least annually, and actions required to achieve this communicated to all parties (patient, carer and GP), with the outcome documented in the patient's notes. The patient's height and weight should be checked and pubertal development if required. Smoking status should be checked in older children. Reports of all outpatient consultations, including the outcome of any drug holidays, should be provided to the GP. Report any suspected adverse drug reactions (ADRs) to the Medicines and Healthcare products Regulatory Agency (MHRA) via the yellow card scheme. When appropriate, undertake periodic treatment withdrawals, or advise the GP in writing how and when to undertake them. Promptly communicate any changes, recommendations, outcomes or other important information to the GP, and provide any other advice or information to the GP or patient if required.
GP Responsibilities	<ul style="list-style-type: none"> Inform secondary care specialist immediately if unable to take on shared care. Prescribe melatonin once the patient is stable on a particular dose, continuing with the formulation selected by the specialist where clinical justification has been provided. Only continue to prescribe melatonin if there is some evidence of ongoing efficacy and benefit. Due to lack of long-term safety data, indefinite continuation of melatonin without specialist review is not recommended. Review the need for continuing melatonin at least every 6 months, referring back to the hospital specialist as appropriate. Seek/follow advice of specialists to facilitate patients having a drug holiday at least annually, and document this and the outcome in patient's notes. Counsel

Commented [SO2]: From Medicines for Children

Commented [SO3]: I've been asked whose responsibility this should be – specialists or GPs? I've assumed that specialists will take responsibility for deciding how and when to do it, but that GPs may need to help out, especially if it's to happen a month before annual review, so have added this sentence. Do specialists concur?

Commented [SO4]: Please can you clarify whether this would be in the hospital notes or GP notes, or both?

Commented [SO5]: Added as received comments that it wasn't clear who should take responsibility for this

Commented [SO6]: Added – do specialists agree?

Commented [SO7]: Added to try and clarify who's doing what!

	<p>patients/carers on the importance of attending outpatient appointments regularly. Do not continue to prescribe melatonin beyond 2 years without specialist review.</p> <ul style="list-style-type: none"> • Ask patient/carer about side effects and general wellbeing, and check smoking status in older patients. Communicate any problems back to the specialist. • If patient needs review, contact the initiating prescriber; CAMHS or paediatric team, but continue to prescribe until the reassessment has taken place (unless an adverse effect has occurred). • Ask the specialist to take back the prescribing should unmanageable problems arise and allow an adequate notice period (4 weeks is a suggested minimum). • Report any suspected adverse drug reactions (ADRs), including any concerns about height, weight or pubertal development, to the specialist and, when appropriate, to the Medicines and Healthcare products Regulatory Agency (MHRA) via the yellow card scheme.
Patient/Carer responsibilities	<ul style="list-style-type: none"> • Report to the specialist or GP if they do not have a clear understanding of their treatment • Contact the GP to arrange supplies of melatonin, allowing sufficient time (usually at least 7 days) to ensure continuity of treatment • Take/give the melatonin as advised. • Attend all booked appointments with hospital and GP, bringing sleep diary (if issued). • Share any concerns in relation to treatment with melatonin with the specialist, GP or pharmacist • Report any side effects or warning symptoms to the specialist, GP or pharmacist whilst taking/giving melatonin. • Report any use of over-the-counter medicines or complementary therapies (e.g. herbal medicines or supplements), illicit drugs and smoking to the specialist, GP or pharmacist whilst taking/giving melatonin.
Disease and drug monitoring	<p>No routine tests required. Specialists should monitor patient's weight and height annually, and pubertal development as required. The patient should be monitored at regular intervals (at least 6 monthly) to check melatonin is still the most appropriate treatment.</p>
Contra-indications	<p>Known hypersensitivity to melatonin or excipients.</p>
Cautions	<p>Refer to BNF or product SmPC/PIL (licensed formulations) published online at www.medicines.org.uk or www.mhra.gov.uk.</p> <ul style="list-style-type: none"> • Melatonin may cause drowsiness. After taking, children should not drive a vehicle, ride a bicycle or use machinery until completely recovered. • Due to lack of clinical data, melatonin is not recommended for use in patients with autoimmune diseases. • Circadin® and Slenyto® tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, the LAPP lactase deficiency or glucose-galactose malabsorption should not take these formulations. • Caution is required in hepatic impairment since the liver is the primary site of melatonin metabolism. • Although caution is advised in product SmPCs, specialist texts advise melatonin can be dosed as normal in all stages of renal impairment.

Commented [SO8]: Do specialists agree?

Commented [SO9]: Added from Slenyto SmPC

Commented [SO10]: added

Commented [SO11]: from Renal Drug Database

Important adverse effects & management	<ul style="list-style-type: none"> • Melatonin is generally well tolerated. Side effects are uncommon. • For a complete list of side effects refer to BNF or product SmPC/PIL (licensed formulations) published online at www.medicines.org.uk or www.mhra.gov.uk. • The most frequently reported adverse effects with melatonin (Slenyto) in children in clinical studies were somnolence, fatigue, mood swings, headache, irritability, aggression and hangover. • Little is known about the long-term effects (>2 years) of melatonin. There are concerns particularly about possible adverse effects on pubertal development in children. Data from clinical studies so far do not support these concerns, but more long term safety data are needed. <p>Report adverse effects to specialist and to the Medicines and Healthcare products Regulatory Agency (MHRA) via the yellow card scheme. If adverse effect is clinically significant, discontinue melatonin and refer patient to specialist.</p>
--	---

4

Prepared by: Kazeem Olalekan, Specialist Pharmacist, Child Health, University Hospital Southampton NHS Foundation Trust and Samantha Owen, Principal Pharmacist, Critical Evaluation and Formulary, University Hospital Southampton NHS Foundation Trust (based on previous guidelines produced by Edward Horne, Pharmacist, Intermediate Care & Community Nursing, Solent NHS Trust, June 2016).

In consultation with: Dr Cathy Hill[and others]

Date prepared: June 2020

Review date: June 2022

Important Interactions	<p>Refer to BNF or product SmPC/PIL (licensed formulations published online at www.medicines.org.uk or www.mhra.gov.uk).</p> <ul style="list-style-type: none"> Fluvoxamine, 5- or 8-methoxypsoralen, cimetidine, oestrogens, and quinolone antibiotics (e.g. ciprofloxacin, norfloxacin) may increase melatonin levels by inhibiting its metabolism resulting in increased risk of adverse effects. Enzyme inducers such as carbamazepine and rifampicin may reduce melatonin levels, requiring dose adjustment. Concomitant administration of other sedatives, e.g. benzodiazepines and "z"- drugs, and tricyclic antidepressants should be avoided due to increased risk of adverse effects. Non-steroidal inflammatory drugs (NSAIDs), e.g. ibuprofen, and beta-blockers may suppress night-time release of endogenous melatonin levels. Administration of these drugs should be avoided in the evening. Alcohol should not be taken with melatonin, because it reduces the effectiveness of melatonin on sleep. Smoking may reduce levels and efficacy of melatonin, while caffeine may increase levels and risk of adverse effects.
-------------------------------	--

Sources of Information/Contacts

University Hospitals Southampton FT	Name / position	Telephone	Email
Specialist / Consultant	Dr Catherine M Hill Children's Sleep Disorder Service Southampton Children's Hospital	Secretary: Leigh Drust 023 8120 5922	Referral forms for the are available via the website: Link here
Hospital Pharmacy	Southampton Medicines Advice Service Kazeem Olalekan Specialist Paediatric Pharmacist	023 8120 6908/9 023 80777 222; Bleep: 1033	medicinesadvice@uhs.nhs.uk kolalekan@nhs.net

Hampshire Hospitals Foundation Trust	Name / position	Telephone	Email
Specialist / Consultant	Dr Rebecca Macleod and Dr Paola Mirto, Consultant Community Paediatricians, Basingstoke and North Hampshire Hospital Dr Rebecca Ferris, Paediatric Consultant, Royal Hampshire County Hospital, Winchester	01256 313083 01962 824995	Paediatric.SecretariesBNH@hhft.nhs.uk Paediatric.SecretariesRHCH@hhft.nhs.uk
Hospital Pharmacy	Helen Leighs, Specialist Paediatric Pharmacist, Basingstoke and North Hampshire Hospital	01256 473202 ext 3341 Bleep: 1560	helen.leighs@hhft.nhs.uk

Sussex Partnership Trust	Name / position	Telephone	Email
--------------------------	-----------------	-----------	-------

Prepared by: Kazeem Olalekan, Specialist Pharmacist, Child Health, University Hospital Southampton NHS Foundation Trust and Samantha Owen, Principal Pharmacist, Critical Evaluation and Formulary, University Hospital Southampton NHS Foundation Trust (based on previous guidelines produced by Edward Horne, Pharmacist, Intermediate Care & Community Nursing, Solent NHS Trust, June 2016).

In consultation with: Dr Cathy Hill[and others]


Date prepared: June 2020

Review date: June 2022

Specialist / Consultant	Havant: Dr Asha Gowda (Consultant, Havant) Dr Fiona Holden (Consultant, Fareham & Gosport)	02392 224560 01329 822220	
Hospital Pharmacy	Worthing Hospital CAMHS Pharmacist	01903 205111 (ext 85698) 07825 118323	pharmacy@wsht.nhs.uk graham.brown@wsht.nhs.uk

Solent NHS Trust	Name / position	Telephone	Email
Consultant	Dr Julia Waine	023 9268 4700	
CAMHS Portsmouth	Falcon House, St James Hospital, Locksway Road, Portsmouth, Hampshire, PO4 9EH	023 9268 4700	SNHS.CAMHS-general@nhs.net
Pharmacy	St Mary's Hospital Portsmouth	023 92680280	

Melatonin Pathway for Children



Prepared by: Ms Ruth Ellenby, Senior Prescribing Support Pharmacist, North Hampshire CCG and Samantha Owen, Principal Pharmacist Critical evaluation, University Hospital Southampton NHS Foundation Trust; March 2020